

Odporúčania Európskej resuscitačnej rady pre kardiopulmonálnu resuscitáciu 2010

Sekcia 2 Základná neodkladná resuscitácia dospelých a používanie automatických externých defibrilátorov

Resuscitation 2010; 81: 1219 - 1276

Zoznam literatúry

1. Recommended guidelines for uniform reporting of data from outofhospital cardiac arrest: the 'Utstein style'. Prepared by a Task Force of Representatives from the European Resuscitation Council, American Heart Association, Heart and Stroke Foundation of Canada, Australian Resuscitation Council. *Resuscitation* 1991;22:1–26.
2. Deakin CD, Nolan JP, Sunde K, Koster RW. European Resuscitation Council Guidelines for Resuscitation 2010. Section 3. Electrical therapies: automated external defibrillators, defibrillation, cardioversion and pacing. *Resuscitation* 2010;81:1293–304.
3. Deakin CD, Nolan JP, Soar J, et al. European Resuscitation Council Guidelines for Resuscitation 2010. Section 4. Adult advanced life support. *Resuscitation* 2010;81:1305–52.
4. Sans S, Kesteloot H, Kromhout D. The burden of cardiovascular diseases mortality in Europe. Task Force of the European Society of Cardiology on Cardiovascular Mortality and Morbidity Statistics in Europe. *Eur Heart J* 1997;18:1231–48.
5. Atwood C, Eisenberg MS, Herlitz J, Rea TD. Incidence of EMStreated outofhospital cardiac arrest in Europe. *Resuscitation* 2005;67:75–80.
6. Cobb LA, Fahrenbruch CE, Olsufka M, Copass MK. Changing incidence of outofhospital ventricular fibrillation, 1980–2000. *JAMA* 2002;288:3008–13.
7. Rea TD, Pearce RM, Raghunathan TE, et al. Incidence of outofhospital cardiac arrest. *Am J Cardiol* 2004;93:1455–60.
8. Vaillancourt C, Verma A, Trickett J, et al. Evaluating the effectiveness of dispatchassisted cardiopulmonary resuscitation instructions. *Acad Emerg Med* 2007;14:877–83.
9. Agarwal DA, Hess EP, Atkinson EJ, White RD. Ventricular fibrillation in Rochester, Minnesota: experience over 18 years. *Resuscitation* 2009;80:1253–8.
10. Ringh M, Herlitz J, Hollenberg J, Rosenqvist M, Svensson L. Out of hospital cardiac arrest outside home in Sweden, change in characteristics, outcome and availability for public access defibrillation. *Scand J Trauma Resusc Emerg Med* 2009;17:18.
11. Cummins R, Thies W. Automated external defibrillators and the Advanced Cardiac Life Support Program: a new initiative from the American Heart Association. *Am J Emerg Med* 1991;9:91–3.
12. Waalewijn RA, Nijpels MA, Tijssen JG, Koster RW. Prevention of deterioration of ventricular fibrillation by basic life support during outofhospital cardiac arrest. *Resuscitation* 2002;54:31–6.
13. Weisfeldt ML, Sitlani CM, Ornato JP, et al. Survival after application of automatic external defibrillators before arrival of the emergency medical system: evaluation in the resuscitation outcomes consortium population of 21 million. *J Am Coll Cardiol* 2010;55:1713–20.
14. van Alem AP, Vrenken RH, de Vos R, Tijssen JG, Koster RW. Use of automated external defibrillator by first responders in out of hospital cardiac arrest: prospective controlled trial. *BMJ* 2003;327:1312.
15. Nolan J, Soar J, Eikeland H. The chain of survival. *Resuscitation* 2006;71:270–1.
16. Muller D, Agrawal R, Arntz HR. How sudden is sudden cardiac death? *Circulation* 2006;114:1146–50.
17. Lowel H, Lewis M, Hormann A. Prognostic significance of prehospital phase in acute myocardial infarct. Results of the Augsburg Myocardial Infarct Registry, 1985–1988. *Dtsch MedWochenschr* 1991;116:729–33.
18. Waalewijn RA, Tijssen JG, Koster RW. Bystander initiated actions in outofhospital cardiopulmonary resuscitation: results from the Amsterdam Resuscitation Study (ARREST). *Resuscitation* 2001;50:273–9.
19. Valenzuela TD, Roe DJ, Cretin S, Spaite DW, Larsen MP. Estimating effectiveness of cardiac arrest interventions: a logistic regression survival model. *Circulation* 1997;96:3308–13.
20. Holmberg M, Holmberg S, Herlitz J. Factors modifying the effect of bystander cardiopulmonary resuscitation on survival in outofhospital cardiac arrest patients in Sweden. *Eur Heart J* 2001;22:511–9.
21. Holmberg M, Holmberg S, Herlitz J, Gardelov B. Survival after cardiac arrest outside hospital in Sweden. Swedish Cardiac Arrest Registry. *Resuscitation* 1998;36:29–36.
22. SOSKANTO Study Group. Cardiopulmonary resuscitation by bystanders with chest compression only (SOSKANTO): an observational study. *Lancet* 2007;369:920–6.

23. Iwami T, Kawamura T, Hiraide A, et al. Effectiveness of bystanderinitiated cardioonly resuscitation for patients with outofhospital cardiac arrest. *Circulation* 2007;116:2900–7.
24. Rea TD, Eisenberg MS, Culley LL, Becker L. Dispatcherassisted cardiopulmonary resuscitation and survival in cardiac arrest. *Circulation* 2001;104:2513–6.
25. Kuisma M, Boyd J, Vayrynen T, Repo J, NousilaWiik M, Holmstrom P. Emergency call processing and survival from outofhospital ventricular fibrillation. *Resuscitation* 2005;67:89–93.
26. Rea TD, Fahrenbruch C, Culley L, et al. CPR with chest compresssions alone or with rescue breathing. *N Engl J Med* 2010;363:423–33.
27. Svensson L, Bohm K, Castren M, et al. Compressiononly CPR or standard CPR in outofhospital cardiac arrest. *N Engl J Med* 2010;363:434–42.
28. Weaver WD, Hill D, Fahrenbruch CE, et al. Use of the automatic external defibrillator in the management of outofhospital cardiac arrest. *N Engl J Med* 1988;319:661–6.
29. Auble TE, Menegazzi JJ, Paris PM. Effect of outofhospital defibrillation by basic life support providers on cardiac arrest mortality: a metaanalysis. *Ann Emerg Med* 1995;25:642–58.
30. Stiell IG, Wells GA, Field BJ, et al. Improved outofhospital cardiac arrest survival through the inexpensive optimization of an existing defibrillation program: OPALS study phase II. Ontario prehospital advanced life support. *JAMA* 1999;281:1175–81.
31. Stiell IG, Wells GA, DeMaio VJ, et al. Modifiable factors associated with improved cardiac arrest survival in a multicenter basic life support/ defibrillation system: OPALS Study Phase I results. Ontario prehospital advanced life support. *Ann Emerg Med* 1999;33:44–50.
32. Caffrey S. Feasibility of public access to defibrillation. *Curr Opin Crit Care* 2002;8:195–8. *R.W. Koster et al. / Resuscitation 81 (2010) 1277–1292* 1291
33. O'Rourke MF, Donaldson E, Geddes JS. An airline cardiac arrest program. *Circulation* 1997;96:2849–53.
34. Page RL, Hamdan MH,McKenas DK. Defibrillation aboard a commercial aircraft. *Circulation* 1998;97:1429–30.
35. Valenzuela TD, Roe DJ, Nichol G, Clark LL, Spaite DW, Hardman RG. Outcomes of rapid defibrillation by security officers after cardiac arrest in casinos. *N Engl J Med* 2000;343:1206–9.
36. Waalewijn RA, de Vos R, Tijssen JG, Koster RW. Survival models for outofhospital cardiopulmonary resuscitation from the perspectives of the bystander, the first responder, and the paramedic. *Resuscitation* 2001;51:113–22.
37. Carr BG, Kahn JM, Merchant RM, Kramer AA, Neumar RW. Interhospital variability in postcardiac arrest mortality. *Resuscitation* 2009;80:30–4.
38. Neumar RW, Nolan JP, Adrie C, et al. Postcardiac arrest syndrome: epidemiology, pathophysiology, treatment, and prognostication. A consensus statement from the International Liaison Committee on Resuscitation (American Heart Association, Australian and New Zealand Council on Resuscitation, European Resuscitation Council, Heart and Stroke Foundation of Canada, InterAmerican Heart Foundation, Resuscitation Council of Asia, and the Resuscitation Council of Southern Africa); the American Heart Association Emergency Cardiovascular Care Committee; the Council on Cardiovascular Surgery and Anesthesia; the Council on Cardiopulmonary, Perioperative, and Critical Care; the Council on Clinical Cardiology; and the Stroke Council. *Circulation* 2008;118:2452–83.
39. Sunde K, Pytte M, Jacobsen D, et al. Implementation of a standardised treatment protocol for post resuscitation care after outofhospital cardiac arrest. *Resuscitation* 2007;73:29–39.
40. Bernard SA, Gray TW, Buist MD, et al. Treatment of comatose survivors of outofhospital cardiac arrest with induced hypothermia. *N Engl J Med* 2002;346:557–63.
41. Mild therapeutic hypothermia to improve the neurologic outcome after cardiac arrest. *N Engl J Med* 2002;346:549–56.
42. Arrich J, Holzer M, Herkner H, Mullner M. Hypothermia for neuroprotection in adults after cardiopulmonary resuscitation. *Cochrane Database Syst Rev* 2009. CD004128.
43. Kitamura T, Iwami T, Kawamura T, Nagao K, Tanaka H, Hiraide A. Nationwide publicaccess defibrillation in Japan. *N Engl J Med* 2010;362:994–1004.
44. Weisfeldt ML, Becker LB. Resuscitation after cardiac arrest: a 3phase timesensitive model. *JAMA* 2002;288:3035–8.
45. White RD, Russell JK. Refibrillation, resuscitation and survival in outofhospital sudden cardiac arrest victims treated with biphasic automated external defibrillators. *Resuscitation* 2002;55:17–23.
46. Kerber RE, Becker LB, Bourland JD, et al. Automatic external defibrillators for public access defibrillation: recommendations for specifying and reporting arrhythmia analysis algorithm performance, incorporating new waveforms, and enhancing safety. A statement for health professionals from the American

Heart Association Task Force on Automatic External Defibrillation, Subcommittee on AED Safety and Efficacy. *Circulation* 1997;95:1677–82.

47. Larsen MP, Eisenberg MS, Cummins RO, Hallstrom AP. Predicting survival from outofhospital cardiac arrest: a graphic model. *Ann Emerg Med* 1993;22:1652–8.

48. Holmberg M, Holmberg S, Herlitz J. Effect of bystander cardiopulmonary resuscitation in outofhospital cardiac arrest patients in Sweden. *Resuscitation* 2000;47:59–70.

49. Aprehmian C, Thompson BM, Finger WA, Darin JC. Experimental cervical spine injury model: evaluation of airway management and splinting techniques. *Ann Emerg Med* 1984;13:584–7.

50. Bahr J, Klingler H, Panzer W, Rode H, Kettler D. Skills of lay people in checking the carotid pulse. *Resuscitation* 1997;35:23–6.

51. Nyman J, Sihvonen M. Cardiopulmonary resuscitation skills in nurses and nursing students. *Resuscitation* 2000;47:179–84.

52. Tibballs J, Russell P. Reliability of pulse palpation by healthcare personnel to diagnose paediatric cardiac arrest. *Resuscitation* 2009;80:61–4.

53. Ruppert M, Reith MW, Widmann JH, et al. Checking for breathing: evaluation of the diagnostic capability of emergency medical services personnel, physicians, medical students, and medical laypersons. *Ann Emerg Med* 1999;34:720–9.

54. Perkins GD, Stephenson B, Hulme J, Monsieurs KG. Birmingham assessment of breathing study (BABS). *Resuscitation* 2005;64:109–13.

55. Hauff SR, Rea TD, Culley LL, Kerry F, Becker L, Eisenberg MS. Factors impeding dispatcher-assisted telephone cardiopulmonary resuscitation. *Ann Emerg Med* 2003;42:731–7.

56. Bobrow BJ, Zuercher M, Ewy GA, et al. Gaspings during cardiac arrest in humans is frequent and associated with improved survival. *Circulation* 2008;118:2550–4.

57. Clark JJ, Larsen MP, Culley LL, Graves JR, Eisenberg MS. Incidence of agonal respirations in sudden cardiac arrest. *Ann Emerg Med* 1992;21:1464–7.

58. Karlsten R, Elowsson P. Who calls for the ambulance: implications for decision support. A descriptive study from a Swedish dispatch centre. *Eur J Emerg Med* 2004;11:125–9.

59. Nurmi J, Pettila V, Biber B, Kuisma M, Komulainen R, Castren M. Effect of protocol compliance to cardiac arrest identification by emergency medical dispatchers. *Resuscitation* 2006;70:463–9.

60. Berdowski J, Beekhuis F, Zwinderman AH, Tijssen JG, Koster RW. Importance of the first link: description and recognition of an outofhospital cardiac arrest in an emergency call. *Circulation* 2009;119:2096–102.

61. Clawson J, Olola C, Heward A, Patterson B. Cardiac arrest predictability in seizure patients based on emergency medical dispatcher identification of previous seizure or epilepsy history. *Resuscitation* 2007;75:298–304.

62. Mithoefer JC, Mead G, Hughes JM, Iliff LD, Campbell EJ. A method of distinguishing death due to cardiac arrest from asphyxia. *Lancet* 1967;2:654–6.

63. Kern KB, Hilwig RW, Berg RA, Sanders AB, Ewy GA. Importance of continuous chest compressions during cardiopulmonary resuscitation: improved outcome during a simulated single layrescuer scenario. *Circulation* 2002;105:645–9.

64. Bobrow BJ, Clark LL, Ewy GA, et al. Minimally interrupted cardiac resuscitation by emergency medical services for outofhospital cardiac arrest. *JAMA* 2008;299:1158–65.

65. Taylor RB, Brown CG, Bridges T, Werman HA, Ashton J, Hamlin RL. A model for regional blood flow measurements during cardiopulmonary resuscitation in a swine model. *Resuscitation* 1988;16:107–18.

66. Aufderheide TP, Sigurdsson G, Pirralo RG, et al. Hyperventilation-induced hypotension during cardiopulmonary resuscitation. *Circulation* 2004;109:1960–5.

67. Eftestol T, Sunde K, Steen PA. Effects of interrupting precordial compressions on the calculated probability of defibrillation success during outofhospital cardiac arrest. *Circulation* 2002;105:2270–3.

68. Wenzel V, Idris AH, Banner MJ, Kubilis PS, Williams JLJ. Influence of tidal volume on the distribution of gas between the lungs and stomach in the nonintubated patient receiving positive-pressure ventilation. *Crit Care Med* 1998;26:364–8.

69. Idris A, Gabrielli A, Caruso L. Smaller tidal volume is safe and effective for bag-valve-ventilation, but not for mouth-to-mouth ventilation: an animal model for basic life support. *Circulation* 1999;100:I–644.

70. Idris A, Wenzel V, Banner MJ, Melker RJ. Smaller tidal volumes minimize gastric inflation during CPR with an unprotected airway. *Circulation* 1995;92(Suppl.):I–759.

71. Dorph E, Wik L, Steen PA. Arterial blood gases with 700 ml tidal volumes during outofhospital CPR. *Resuscitation* 2004;61:23–7.

72. Winkler M, Mauritz W, Hackl W, et al. Effects of half the tidal volume during cardiopulmonary resuscitation on acid–base balance and haemodynamics in pigs. *Eur J Emerg Med* 1998;5:201–6.

73. Ruben H. The immediate treatment of respiratory failure. *Br J Anaesth* 1964;36:542–9.
74. Elam JO. Bagvalvemask O₂ ventilation. In: Safar P, Elam JO, editors. *Advances in cardiopulmonary resuscitation: the Wolf Creek conference on cardiopulmonary resuscitation*. New York, NY: SpringerVerlag, Inc.; 1977. p. 73–9.
75. Dailey RH. *The airway: emergency management*. St. Louis, MO: Mosby Year Book; 1992.
76. Paradis NA, Martin GB, Goetting MG, et al. Simultaneous aortic, jugular bulb, and right atrial pressures during cardiopulmonary resuscitation in humans. Insights into mechanisms. *Circulation* 1989;80:361–8.
77. KramerJohansen J, Myklebust H, Wik L, et al. Quality of outofhospital cardiopulmonary resuscitation with real time automated feedback: a prospective interventional study. *Resuscitation* 2006;71:283–92.
78. Edelson DP, Abella BS, KramerJohansen J, et al. Effects of compression depth and preshock pauses predict defibrillation failure during cardiac arrest. *Resuscitation* 2006;71:137–45.
79. Wik L, KramerJohansen J, Myklebust H, et al. Quality of cardiopulmonary resuscitation during outofhospital cardiac arrest. *JAMA* 2005;293:299–304.
80. Abella BS, Alvarado JP, Myklebust H, et al. Quality of cardiopulmonary resuscitation during inhospital cardiac arrest. *JAMA* 2005;293:305–10.
81. Christenson J, Andrusiek D, EversonStewart S, et al. Chest compression fraction determines survival in patients with outofhospital ventricular fibrillation. *Circulation* 2009;120:1241–7.
82. Ochoa FJ, RamalleGomara E, Carpintero JM, Garcia A, Saralegui I. Competence of health professionals to check the carotid pulse. *Resuscitation* 1998;37:173–5.
83. Shin J, Rhee JE, Kim K. Is the internipple line the correct hand position for effective chest compression in adult cardiopulmonary resuscitation? *Resuscitation* 2007;75:305–10.
84. Kusunoki S, Tanigawa K, Kondo T, Kawamoto M, Yuge O. Safety of the internipple line hand position landmark for chest compression. *Resuscitation* 2009;80:1175–80.
85. Delvaux AB, Trombley MT, Rivet CJ, et al. Design and development of a cardiopulmonary resuscitation mattress. *J Intensive Care Med* 2009;24:195–9.
86. Perkins GD, Smith CM, Augre C, et al. Effects of a backboard, bed height, and operator position on compression depth during simulated resuscitation. *Intensive Care Med* 2006;32:1632–5.
87. Perkins GD, Kocierz L, Smith SC, McCulloch RA, Davies RP. Compression feedback devices over estimate chest compression depth when performed on a bed. *Resuscitation* 2009;80:79–82.
88. Aufderheide TP, Pirralo RG, Yannopoulos D, et al. Incomplete chest wall decompression: a clinical evaluation of CPR performance by EMS personnel and assessment of alternative manual chest compression–decompression techniques. *Resuscitation* 2005;64:353–62.
89. Yannopoulos D, McKnite S, Aufderheide TP, et al. Effects of incomplete chest wall decompression during cardiopulmonary resuscitation on coronary and cerebral perfusion pressures in a porcine model of cardiac arrest. *Resuscitation* 2005;64:363–72.
90. Sanders AB, Kern KB, Berg RA, Hilwig RW, Heidenrich J, Ewy GA. Survival and neurologic outcome after cardiopulmonary resuscitation with four different chest compression–ventilation ratios. *Ann Emerg Med* 2002;40:553–62.
91. Dorph E, Wik L, Stromme TA, Eriksen M, Steen PA. Quality of CPR with three different ventilation:compression ratios. *Resuscitation* 2003;58:193–201.
92. Dorph E, WikL, Stromme TA, Eriksen M, Steen PA. Oxygen delivery and return of spontaneous circulation with ventilation:compression ratio 2:30 versus chest compressions only CPR in pigs. *Resuscitation* 2004;60:309–18. 1292 *R.W. Koster et al. / Resuscitation 81 (2010) 1277–1292*
93. Babbs CF, Kern KB. Optimum compression to ventilation ratios in CPR under realistic, practical conditions: a physiological and mathematical analysis. *Resuscitation* 2002;54:147–57.
94. Fenici P, Idris AH, Lurie KG, Ursella S, Gabrielli A. What is the optimal chest compression–ventilation ratio? *Curr Opin Crit Care* 2005;11:204–11.
95. Sayre MR, Cantrell SA, White LJ, Hiestand BC, Keseg DP, Koser S. Impact of the 2005 American Heart Association cardiopulmonary resuscitation and emergency cardiovascular care guidelines on outofhospital cardiac arrest survival. *Prehosp Emerg Care* 2009;13:469–77.
96. Olasveengen TM, Vik E, Kuzovlev A, Sunde K. Effect of implementation of new resuscitation guidelines on quality of cardiopulmonary resuscitation and survival. *Resuscitation* 2009;80:407–11.
97. Aufderheide TP, Lurie KG. Death by hyperventilation: a common and lifethreatening problem during cardiopulmonary resuscitation. *Crit Care Med* 2004;32:S345–51.
98. Ornato JP, Hallagan LF, McMahan SB, Peeples EH, Rostafinski AG. Attitudes of BCLS instructors about mouthto mouth resuscitation during the AIDS epidemic. *Ann Emerg Med* 1990;19:151–6.
99. Hew P, Brenner B, Kaufman J. Reluctance of paramedics and emergency medical technicians to perform mouthto mouth resuscitation. *J Emerg Med* 1997;15:279–84.

100. Chandra NC, Gruben KG, Tsitlik JE, et al. Observations of ventilation during resuscitation in a canine model. *Circulation* 1994;90:3070–5.
101. Geddes LA, Rundell A, Otlewski M, Pargett M. How much lung ventilation is obtained with only chestcompression CPR? *Cardiovasc Eng* 2008;8:145–8.
102. Berg RA, Kern KB, Hilwig RW, et al. Assisted ventilation does not improve outcome in a porcine model of singlerescuer bystander cardiopulmonary resuscitation. *Circulation* 1997;95:1635–41.
103. Berg RA, Kern KB, Hilwig RW, Ewy GA. Assisted ventilation during ‘bystander’ CPR in a swine acute myocardial infarction model does not improve outcome. *Circulation* 1997;96:4364–71.
104. Turner I, Turner S, Armstrong V. Does the compression to ventilation ratio affect the quality of CPR: a simulation study. *Resuscitation* 2002;52:55–62.
105. Bohm K, Rosenqvist M, Herlitz J, Hollenberg J, Svensson L. Survival is similar after standard treatment and chest compression only in outofhospital bystander cardiopulmonary resuscitation. *Circulation* 2007;116:2908–12.
106. Kitamura T, Iwami T, Kawamura T, Nagao K, Tanaka H, Hiraide A. Bystanderinitiated rescue breathing for outofhospital cardiac arrests of noncardiac origin. *Circulation* 2010;122:293–9.
107. Kitamura T, Iwami T, Kawamura T, et al. Conventional and chestcompressiononly cardiopulmonary resuscitation by bystanders for children who have outofhospital cardiac arrests: a prospective, nationwide, populationbased cohort study. *Lancet* 2010;375:1347–54.
108. Handley AJ, Handley JA. Performing chest compressions in a confined space. *Resuscitation* 2004;61:55–61.
109. Perkins GD, Stephenson BT, Smith CM, Gao F. A comparison between overhead and standard cardiopulmonary resuscitation. *Resuscitation* 2004;61:155–61.
110. White L, Rogers J, Bloomingdale M, et al. Dispatcherassisted cardiopulmonary resuscitation: risks for patients not in cardiac arrest. *Circulation* 2010;121:91–7.
111. Cheung W, Gullick J, Thanakrishnan G, et al. Injuries occurring in hospital staff attending medical emergency team (MET) calls—a prospective, observational study. *Resuscitation* 2009;80:1351–6.
112. Sullivan F, Avstreich D. Pneumothorax during CPR training: case report and review of the CPR literature. *Prehosp Disaster Med* 2000;15:64–9.
113. Peberdy MA, Ottingham LV, Groh WJ, et al. Adverse events associated with lay emergency response programs: the public access defibrillation trial experience. *Resuscitation* 2006;70:59–65.
114. Sugerma NT, Edelson DP, Leary M, et al. Rescuer fatigue during actual inhospital cardiopulmonary resuscitation with audiovisual feedback: a prospective multicenter study. *Resuscitation* 2009;80:981–4.
115. Hallstrom AP, Ornato JP, Weisfeldt M, et al. Publicaccess defibrillation and survival after outofhospital cardiac arrest. *N Engl J Med* 2004;351:637–46.
116. Hoke RS, Heinroth K, Trappe HJ, Werdan K. Is external defibrillation an electric threat for bystanders? *Resuscitation* 2009;80:395–401.
117. Axelsson A, Herlitz J, Karlsson T, et al. Factors surrounding cardiopulmonary resuscitation influencing bystanders’ psychological reactions. *Resuscitation* 1998;37:13–20.
118. Axelsson A, Herlitz J, Ekstrom L, Holmberg S. Bystanderinitiated cardiopulmonary resuscitation outofhospital. A first description of the bystanders and their experiences. *Resuscitation* 1996;33:3–11.
119. Mejicano GC, Maki DG. Infections acquired during cardiopulmonary resuscitation: estimating the risk and defining strategies for prevention. *Ann Intern Med* 1998;129:813–28.
120. Cydulka RK, Connor PJ, Myers TF, Pavza G, Parker M. Prevention of oral bacterial flora transmission by using mouthtomask ventilation during CPR. *J Emerg Med* 1991;9:317–21.
121. Blenkharn JI, Buckingham SE, Zideman DA. Prevention of transmission of infection during mouthtomouth resuscitation. *Resuscitation* 1990;19:151–7.
122. Turner S, Turner I, Chapman D, et al. A comparative study of the 1992 and 1997 recovery positions for use in the UK. *Resuscitation* 1998;39:153–60.
123. Handley AJ. Recovery Position. *Resuscitation* 1993;26:93–5.
124. Anonymous. Guidelines 2000 for Cardiopulmonary resuscitation and emergency cardiovascular care—an international consensus on science. *Resuscitation* 2000;46:1–447.
125. Fingerhut LA, Cox CS, Warner M. International comparative analysis of injury mortality. Findings from the ICE on injury statistics. *International Collaborative Effort on Injury Statistics. Adv Data* 1998:1–20.
126. Proceedings of the 2005 International Consensus on Cardiopulmonary Resuscitation and Emergency Cardiovascular Care Science with Treatment Recommendations. *Resuscitation* 2005;67:157–341.
127. Redding JS. The choking controversy: critique of evidence on the Heimlich maneuver. *Crit Care Med* 1979;7:475–9.
128. Langhelle A, Sunde K, Wik L, Steen PA. Airway pressure with chest compressions versus Heimlich manoeuvre in recently dead adults with complete airway obstruction. *Resuscitation* 2000;44:105–8.

129. Guildner CW, Williams D, Subitch T. Airway obstructed by foreign material: the Heimlich maneuver. *JACEP* 1976;5:675–7.
130. Ruben H, Macnaughton FI. The treatment of foodchoking. *Practitioner* 1978;221:725–9.
131. Hartrey R, Bingham RM. Pharyngeal trauma as a result of blind finger sweeps in the choking child. *J Accid Emerg Med* 1995;12:52–4.
132. Elam JO, Ruben AM, Greene DG. Resuscitation of drowning victims. *JAMA* 1960;174:13–6.
133. Ruben HM, Elam JO, Ruben AM, Greene DG. Investigation of upper airway problems in resuscitation, 1: studies of pharyngeal xrays and performance by laymen. *Anesthesiology* 1961;22:271–9.
134. Kabbani M, Goodwin SR. Traumatic epiglottis following blind finger sweep to remove a pharyngeal foreign body. *Clin Pediatr (Phila)* 1995;34:495–7.
- 134a. European Resuscitation Council Guidelines for Resuscitation 2010: Section 6: Paediatric life support. *Resuscitation* 2010; 81:1400–33.
- 134b. European Resuscitation Council Guidelines for Resuscitation 2010: Section 8: Cardiac arrest in special circumstances. *Resuscitation* 2010; 81:1364–88.
135. Atkins DL, EversonStewart S, Sears GK, et al. Epidemiology and outcomes from outofhospital cardiac arrest in children: the Resuscitation Outcomes Consortium EpistryCardiac Arrest. *Circulation* 2009;119:1484–91.
136. BarCohen Y, Walsh EP, Love BA, Cecchin F. First appropriate use of automated external defibrillator in an infant. *Resuscitation* 2005;67:135–7.
137. Divekar A, Soni R. Successful parental use of an automated external defibrillator for an infant with longQT syndrome. *Pediatrics* 2006;118:e526–9.
138. RodriguezNunez A, LopezHerce J, Garcia C, Dominguez P, Carrillo A, Bellon JM. Pediatric defibrillation after cardiac arrest: initial response and outcome. *Crit Care* 2006;10:R113.
139. Samson RA, Nadkarni VM, Meaney PA, Carey SM, Berg MD, Berg RA. Outcomes of inhospital ventricular fibrillation in children. *N Engl J Med* 2006;354:2328–39.
140. Cobb LA, Fahrenbruch CE, Walsh TR, et al. Influence of cardiopulmonary resuscitation prior to defibrillation in patients with outofhospital ventricular fibrillation. *JAMA* 1999;281:1182–8.
141. Wik L, Hansen TB, Fylling F, et al. Delaying defibrillation to give basic cardiopulmonary resuscitation to patients with outofhospital ventricular fibrillation: a randomized trial. *JAMA* 2003;289:1389–95.
142. Jacobs IG, Finn JC, Oxer HF, Jelinek GA. CPR before defibrillation in outofhospital cardiac arrest: a randomized trial. *Emerg Med Australas* 2005;17:39–45.
143. Baker PW, Conway J, Cotton C, et al. Defibrillation or cardiopulmonary resuscitation first for patients with outofhospital cardiac arrests found by paramedics to be in ventricular fibrillation? A randomised control trial. *Resuscitation* 2008;79:424–31.
144. Indik JH, Hilwig RW, Zuercher M, Kern KB, Berg MD, Berg RA. Preshock cardiopulmonary resuscitation worsens outcome from circulatory phase ventricular fibrillation with acute coronary artery obstruction in swine. *Circ Arrhythm Electrophysiol* 2009;2:179–84.
145. Monsieurs KG, Vogels C, Bossaert LL, Meert P, Calle PA. A study comparing the usability of fully automatic versus semiautomatic defibrillation by untrained nursing students. *Resuscitation* 2005;64:41–7.
146. White RD, Bunch TJ, Hankins DG. Evolution of a communitywide early defibrillation programme experience over 13 years using police/fire personnel and paramedics as responders. *Resuscitation* 2005;65:279–83.
147. Mosesso Jr VN, Davis EA, Auble TE, Paris PM, Yealy DM. Use of automated external defibrillators by police officers for treatment of outofhospital cardiac arrest. *Ann Emerg Med* 1998;32:200–7.
148. The public access defibrillation trial investigators. Publicaccess defibrillation and survival after outofhospital cardiac arrest. *N Engl J Med* 2004;351:637–46.
149. Priori SG, Bossaert LL, Chamberlain DA, et al. Policy statement: ESC–ERC recommendations for the use of automated external defibrillators (AEDs) in Europe. *Resuscitation* 2004;60:245–52.
150. Groh WJ, Newman MM, Beal PE, Fineberg NS, Zipes DP. Limited response to cardiac arrest by police equipped with automated external defibrillators: lack of survival benefit in suburban and rural Indiana—the police as responder automated defibrillation evaluation (PARADE). *Acad Emerg Med* 2001;8:324–30.
151. Sayre MR, Swor R, Pepe PE, Overton J. Current issues in cardiopulmonary resuscitation. *Prehosp Emerg Care* 2003;7:24–30.
152. Nichol G, Hallstrom AP, Ornato JP, et al. Potential costeffectiveness of public access defibrillation in the United States. *Circulation* 1998;97:1315–20.
153. Nichol G, Valenzuela T, Roe D, Clark L, Huszti E, Wells GA. Cost effectiveness of defibrillation by targeted responders in public settings. *Circulation* 2003;108:697–703.

154. Bardy GH, Lee KL, Mark DB, et al. Home use of automated external defibrillators for sudden cardiac arrest. *N Engl J Med* 2008;358:1793–804